




UNIVERSITY OF AGRONOMIC SCIENCES
AND VETERINARY MEDICINE OF BUCHAREST
FACULTY OF ANIMAL SCIENCE



SCIENTIFIC PAPERS

SERIES D. ANIMAL SCIENCE

VOLUME LV



2012
BUCHAREST

DETERMINATION OF ROMANIAN ALFALFA CRUDE PROTEIN AND CRUDE FIBER CONTENTS AS WELL AS *IN VITRO* ORGANIC MATTER DIGESTIBILITY BY NIR SPECTROMETRY

Laura M. DALE^{1,2}, André THEWIS¹, Ioan ROTAR², Christelle BOUDRY¹, Roxana VIDICAN², Anamaria MALINAS², Vasile FLORIAN³, Bernard LECLER⁴, Richard AGNEESSENS⁵, Juan A. FERNÁNDEZ PIERNA⁴, Vincent BAETEN⁴

¹ Animal Science Unit, Gembloux Agro-Bio Tech, University of Liège, 2, Passage des Déportés, 5030 Gembloux, Belgium, E-mail: dale_lm@yahoo.com; dale.laura@student.ulg.ac.be; athewis@ulg.ac.be; christelle.boudry@ulg.ac.be

² Department of Grassland and Forage Crops, University of Agricultural Science and Veterinary Medicine Cluj, 3-5, Calea Mănăştur, 400372, Cluj-Napoca, E-mail: rotarioan52@yahoo.fr; roxanavidican@yahoo.com; ciure_ana@yahoo.com;

³ Department of Phitophatology, University of Agricultural Science and Veterinary Medicine Cluj, 3-5, Calea Mănăştur, 400372, Cluj Napoca, E-mail: vasileflorian2003@yahoo.com

⁴ Walloon Agricultural Research Centre, Valorisation of Agricultural Products Department, 24 Chaussée de Namur, 5030 Gembloux, Belgium, E-mail: fernandez@cra.wallonie.be; baeten@cra.wallonie.be; lecler@cra.wallonie.be.

⁵ Walloon Agricultural Research Centre, Valorisation of Agricultural Products Department, 100 Rue du Serpont, 6800 Libramont-Chevigny, Belgium, E-mail: agnessens@cra.wallonie.be

Corresponding author email: dale_lm@yahoo.com

Abstract

Alfalfa (Medicago sativa L.) is a high quality forage which has been used worldwide. The superiority of alfalfa lies in its high yield, high protein content and high digestibility. The aim of this study was to develop a simple, fast and non-destructive method, named Near Infrared Spectroscopy (NIRS) to determinate alfalfa quality. To realize this study, alfalfa samples were obtained from Mănăştur Experimental Station – Farm Cojocna in 2008–2009, in one experiment carried out using randomization blocks design with two experimental factors (mineral fertilization and period of harvest). Alfalfa quality was first determined on 48 samples by classical analyses: crude protein CP (AOAC, 1990), crude fiber CF (Fiber Cap, FOSS, DK) and in vitro organic matter digestibility OMDrt (DeBoever, 1986). Then the samples were scanned by NIRS. Calibration models were performed by PerkinElmer Spectrum Quant + 4.21 program (USA) on the 48 samples determined by classical analysis. The results showed fully confirmed by acceptable coefficients of determination and standard error of cross-validation ($R^2=0.96$ for CP, 0.94 for CF, 0.98 for OMDrt and $SECV=0.77$ for CP, 1.35 for CF, 1.13 for OMDrt). Successful results for prediction of other 176 alfalfa samples were then obtained using these calibration models: $SEP=0.869$ for CP, 1.058 for CF, 1.058 for OMDrt). The highest CP and OMDrt values of alfalfa were obtained in bud stage (22.0% and 66.0% respectively). While for CF, the highest content was registered in the seed formation stage (46.0%). The NIRS technique offers us the possibility to determinate rapidly and easily Romanian alfalfa important parameters, but the system could also be used for the determination of other constituents.

Keywords: alfalfa, NIRS, crude protein, crude fiber, in vitro organic matter digestibility

INTRODUCTION

In current agriculture, forage production, obtained from permanent grasslands, temporary grasslands and forage crops is an integral part of agricultural land management [1]. A correct appreciation of feed quality requires a set of general analyses such as: botanical composition, organic and mineral contents, palatability and organic matter digestibility, as well as dry matter intake [2].

Alfalfa (*Medicago sativa* L.) is a high quality forage used worldwide. The superiority of alfalfa lies in its high yield, high protein content and high digestibility. It is considered by researchers as the 'queen of forages' [3]. Archaeological informations and antic scientific papers pointed out that alfalfa was cultivated first in South-West of Asia already on 4000 years before Christ [4]. In Romania, alfalfa was first cultivated in the last part of XVIIIth century in Transylvania and Banat

Country [4]. On the agrobiologic point of view alfalfa presents some particularities such as: water resistance, low temperature resistance, good utilization of irrigation water, high capacity of regeneration after harvest period and high competitive supply [3]. Alfalfa has also a high economic importance, manifested by its high ecological plasticity and high productivity. In natural conditions, in Romania alfalfa can offers more than 50 t/ha of green mass on 3 harvests/year and in irrigation conditions more than 80 t/ha of green mass on 4 harvests/year [3]. Jarrige, [5] pointed out that alfalfa has lower proportion of leaves in the first stage of vegetation in comparison with gramineae. But advancing in vegetation stages, alfalfa crude protein content and digestibility value decrease while cell wall content increases. The main components of cell walls are fibers. The organic matter digestibility of the feed depends mainly on the fiber content and its digestibility. Digestibility decreases with the proportion of cell walls and their level of lignification in the plant. Organic matter digestibility of the feed during the first cycle of vegetation depends almost exclusively on the stage of development. Organic matter digestibility of the feed (~60%) at accrued harvest stages has the same significant high value, no matter the place, year and the level of nitrogen fertilization [5]. Routine analyses of forages are necessary to evaluate their nutritive values and calculate optimal and well balanced diets for ruminants. NIRS is a non-destructive method of analysis based on diffuse reflectance of ground samples. NIRS was applied first in the agricultural sector by Norris in the 1960's to determine moisture content in soybeans [6]. More recently, Dale et al. [7] reviewed a lot of studies regarding the use of NIRS and NIR-HSI. This technology was used particularly for fast determination of the nutrients' concentration and feeding value of dried and fresh crop materials [8-11]. NIRS technique has become a 'clean technology' very used in sustainable agriculture [10] because it is using a reduced amount of chemical substances (4, 12-13). The aim of this study was to develop a simple, fast and non-destructive method to determine alfalfa quality (crude protein CP, crude fiber CF and *in vitro* organic matter

digestibility OMDrt), based on Near Infrared Spectroscopy (NIRS system).

MATERIAL AND METHOD

Alfalfa samples were obtained from Mănăstur Experimental Station – Farm Cojocna in 2008–2009, in one experiment carried out using block randomization design with two experimental factors (mineral fertilization and period of harvest). The experiment started in spring 2007, with the seeding of alfalfa. The mineral fertilization was applied each year in early spring in different levels of graduation: unfertilized or fertilized with N₅₀, N₅₀P₅₀, N₅₀P₅₀K₅₀, N₁₀₀, N₁₀₀P₇₅ or N₁₀₀P₇₅K₇₅. The samples were collected directly in the field in the harvest period (plant height 30 cm, bud, flowering and seed formation), then were dried at 60°C in oven 2 days. In the next step, the samples were milled using a 5 mm screen (Grindomix GM 200, Retsch, Haan, Germany) and then milled through a 1 mm sieve (Cyclotec™ 1093, Tecator, Sweden).

Alfalfa quality was determinated on 48 samples by classical analyses: crude protein (CP) (Kjeldhal method [14]), crude fiber (CF) (Weende Scheme - Fiber Cap, FOSS, DK) and *in vitro* organic matter digestibility (OMDrt) (DeBoever method [15]). The same 48 samples were scanned for NIRS analyses to perform calibration model (36 samples) and validation model (12 samples); other 176 samples from the same experiment were then scanned for prediction. Spectra were collected over the wavelengths range 10000 to 4000 cm⁻¹ (1000-2500 nm) at 8 cm⁻¹ resolution with 2 cm⁻¹ step, in two repetitions. Each spectrum is a mean of 32 scans of the same sample.

For the mathematical models, the program PerkinElmer Spectrum Quant + 4.21 was used. The algorithm used for calibration models was Partial Least Squares (PLS). To perform a robust calibration model, spectra were treated by following mathematical treatment: 2nd Derivative (order: 3, window: 15 pt) and a cross validation was also used: leave one out, using 18 PLS factors.

RESULTS AND DISCUSSIONS

The CP, CF and OMDrt contents of alfalfa determined by classical methods are presented in table 1. The results were similar to results showed by Varga [12], Vintilă [16] and

Vîntu et al., [13]. And it can be pointed out those higher levels of crude protein content in the first and second harvested stages where obtained because of a great proportion of leaves than stems in these two stages compared to flowering and seed formation [17].

Table 1. Crude protein (CP), crude fiber (CF) and in vitro organic matter digestibility (OMDrt) of alfalfa harvested at different stages determined by classical methods

Harvest stage	CP	CF	OMDrt
Plant height 30 cm	14.23% - 19.79%	27.92% - 34.37%	57.30% - 63.66%
Bud	14.38% - 17.64%	30.33% - 33.75%	63.39% - 65.83%
Flowering	12.87% - 17.44%	32.03% - 39.82%	52.95% - 59.86%
Seed formation	12.53% - 15.19%	38.05% - 44.03%	38.05% - 44.03%

The PLS algorithm for calibration model was performed on the spectra with the help of data obtained by classical analyses. The characteristics of alfalfa calibration models for CP, CF and OMDrt are presented in table 2. The report SD/SECV allows a comparison of the equations developed, which are independent of units and have been used by other authors. The ratio SD/SECV of 2.5-3.0 has been considered appropriate for measuring samples quality, but to ensure the models robustness, values of at least 3.0-5.0 are necessary [18].

The SD/SECV ratio was 4.18, 3.97 6.70 for CP, CF and OMDrt respectively. The ratios for CP and CF were in the limits presented by Williams and Sobering, [18]. But the ratio for OMDrt was higher than the limits; this means that it will be necessary to perform more classical analyses for the OMDrt model. The results obtained for calibration and validation models were similar to those of other authors for the same type of biological material (table 3).

Table 2. Characteristics of alfalfa calibration models

Content	CP	CF	OMDrt
N	48	48	48
Mean	15.95	27.67	64.7
SEC	0.72	1.32	0.99
R ²	0.96	0.94	0.98
SECV	0.77	1.35	1.13
SEP	0.869	1.058	1.058
SD	3.22	5.37	7.58
SD/SECV	4.18	3.97	6.70

Legend: N: samples taken in calibration model, Mean: the mean value, SEC: the standard error of calibration, R²: the coefficient of determinations, SECV: the standard error for cross validation, SEP: the standard error of prediction, SD: the standard deviation.

Table 3. The coefficient of multiple determinations (R²) and the standard error for cross validation (SECV) of alfalfa calibration models

Calibration models for	R ²	SECV	Reference
CP	0.96	0.96	Sheaffer, et al., [19];
	0.95	0.94	Velasco, et al., [20];
	0.92	0.91	Iantcheva et al., [21];
	0.95	0.94	Brogna et al., [22].
CF	0.87	-	Brogna et al., [22];
	-	3.12.	Iantcheva et al., [21].
OMDrt	-	2.97	Iantcheva et al., [21];
	0.87	3.70	Brogna et al., [22].

Successful results for prediction of other 176 alfalfa samples were also obtained by NIRS analysis (SEP=0.869 for CP, 1.058 for CF, 1.058 for OMDrt - table 1). This system offers

us the possibility of using NIRS technique for alfalfa organic substances determination. The accuracy and reliability of the prediction of e.g. the CP, CF or OMDrt of a sample by this NIRS technique has been totally dependent on the

accuracy and reliability of the classical determination of CP, CF or OMDrt. It is the reason why it is absolutely necessary to

perform accurate analyses by classical techniques.

Table 4. Crude protein (CP), crude fiber (CF) and *in vitro* organic matter digestibility (OMDrt) of alfalfa harvested at different harvest stage determined by NIR spectrometry

Harvest stage	CP	CF	OMDrt
Plant height 30 cm	12.83% - 20.61%	15.96% - 33.61%	50.07% - 63.57%
Bud	14.32% - 22.04%	19.53% - 35.06%	46.46% - 65.99%
Flowering	12.06% - 20.14%	17.56% - 38.47%	45.07% - 61.58%
Seed formation	11.47% - 16.83%	18.33% - 46.06%	33.16% - 45.14%

The *in vitro* OMDrt values of alfalfa in harvest stage were indirectly related to the CF content and directly proportional to the CP content. During the years of the experience, 2008 and 2009, CP levels and OMDrt were the highest at bud stage. A lower OMDrt and a highest CF content were obtained in seed formation stage, showing the role of CF in the OMDrt; the

higher the crude fiber content is, the lower the digestibility is. It can be seen in table 5 that our results obtained by NIRS method were similar to different results obtained by classical or NIRS analysis. The results are presented in function of the harvest stage (bud, flowering and seed formation).

Table 5. Crude protein (CP), crude fiber (CF) and *in vitro* organic matter digestibility (OMDrt) of alfalfa harvested at different stages

Harvest stage	CP	CF	OMDrt
Bud	20.07% [12]; 24.05% [13]; 19.80% [16]; 22.60% [19]; 19.90% [23]; 16.00% [24]; 23.03% [25].	28.70% [12]; 21.40% [13]; 33.40% [24]; 25.49% [25].	50.09% [23]; 63.00% [24].
Flowering	18.47% [12]; 21.17% [13]; 18.10% [16]; 19.20% [19]; 19.40% [23]; 13.20% [24]; 33.09% [25].	31.03% [12]; 28.50% [13]; 35.70% [24]; 29.66% [25].	51.20% [23]; 62.00% [24].
Seed formation	15.92% [12]; 16.45% [13]; 17.50% [16]; 16.40% [19]; 17.40% [23]; 12.30% [24]; 16.37% [25].	34.46% [12]; 32.00% [13]; 33.60% [24]; 40.18% [25].	45.40% [23]; 60.00% [24].

In the first year of experiment a lower content of crude protein was obtained compared to second year for the different harvest stages, because in the first year of the experiment the alfalfa is installing in the yield and the nitrogen is used for this [3]. The same effect was pointed out by Decruyenaere et al., [11] and Stanacev et al., [26]. According to the harvest stage it was observed a reduction of CP and an increase in CF in the latest harvest which could be explained by to the evolution of stems and leaves [17] leaves containing more CP and less CF than stems. Moreover Heinrichs [27] and Babinec et al. [28] pointed out that losses of leaves are important because the protein concentration was higher in leaves than in stems. The crude protein content and the crude fiber contents vary between very wide limits,

depending largely on the development stage of alfalfa [1].

The relative contribution of qualitative parameters mentioned above helps to lift organic matter digestibility up to 60%, contributing equally to protein and fiber content. From this it can be concluded that the digestibility and crude protein content decrease quickly, while crude fiber increase slightly and then remain constant. Demarquilly and Andrieu [17] noted that the whole plant digestibility of alfalfa has been correlated to the proportion of leaves and stems, because leaves are more digestible than stems.

CONCLUSIONS

Based on the samples supplied, it has been shown that NIRS and PLS can be used to determine CP, CF and *in vitro* OMD_rt of alfalfa.

This preliminary study proves that NIRS is an extremely reliable, non-destructive and rapid technique for the prediction of quantitative chemical and physical properties of alfalfa from Romania.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Gembloux Agro Bio-Tech, University of Liege and USAMV Cluj Napoca for their financial support, and the Walloon Agricultural Research Centre in Gembloux, Belgium, for scientific support and advice.

REFERENCES

- [1] Dale L. M., 2011. *Determination of forage quality by destructive and non-destructive methods*. PhD thesis. USAMV Cluj, 330p.
- [2] Rotar, I., Carlier, L., 2011. *Cultura pajistilor*. Cluj - Napoca: Ed. Risoprint, 408p.
- [3] Rotar, I., 1993. *Cercetări privind coacțiile de competiție în culturi pure și asociate de Medicago sativa și Dactylis glomerata în condiții de fertilizare diferențiată cu azot*. Cluj - Napoca: Doctoral Thesis, 230p.
- [4] Florian, C. V., 2009. *Studiul influenței principalelor boli ale lucernei asupra recoltei și calității furajului*. Cluj-Napoca: Doctoral Thesis, 250p.
- [5] Jarrige, R., 1988. *Alimentation des bovins, ovins & caprins*. Paris: INRA.
- [6] Hart, J., Norris, K., Golumbie, C., 1962. *Determination of the moisture content of seeds by near-infrared spectroscopy*. Cereal Chem., 39: 94-99.
- [7] Dale, L. M., Théwis, A., Boudry, C., Rotar, I., Dardenne, P., Baeten, V., Fernández Pierna, J. A., 2012. *Hyperspectral imaging applications in agriculture and agro-food product quality and safety control: A review*. Appl. Spectrosc. Rev. (submitted and accepted article).
- [8] Biston, R., Dardenne, P., Demarquilly, C., 1989. *Determination of forage in vivo digestibility by NIRS*. Proceedings of the XVI International Grassland Congress, Association Française pour la Production Fourragère Versailles (Editor). Nice, France: 895-896.
- [9] Dardenne, P., Agneessens, R., Biston, R., 1992. *La spectrométrie dans le proche infrarouge, outil analytique de gestion de la qualité des produits agricoles et de l'environnement*. Bull. des Rech. Agron. Gembloux, 27: 39-51.
- [10] Vidican, R., Rotar, I., Sima, N., 2000. *Tehnica NIRS (Near Infrared Reflectance Spectroscopy) și aplicațiile sale în analiza calității furajelor*. Simpozion "Agriultura și alimentația", USAMV Cluj Napoca, 187-191.
- [11] Decruyenaere, V., Lecomte, P., Demarquilly, C., Aufrere, J., Dardenne, P., Stilmant, D., et al., 2008. *Evaluation of green forage intake and digestibility in ruminants using near infrared reflectance spectroscopy (NIRS): Developing a global calibration*. Animal Feed Science and Technology, 148, 138-156.
- [12] Varga, P., Moga, I., Kellner, E., Bălan, C., Ionescu, M., 1973. *Lucerna*. București: Ceres, 301p.
- [13] Vintu, V., Stavarache, M., Samuil, C., Muntean, I., 2012. *Chemical Composition dynamics of alfalfa (Medicago sativa L.) at different plant growth stages*. EGF 2012, Grassland Science in Europe, 17, 394-396.
- [14] AOAC, 1990. *Association of Official Analytical Chemists, Methods of Analysis* 15th Edition. Washington D.C.
- [15] De Boever, J., Cottyn, B.G., Buysse, F.X., Wainman, F.W., Vanacker, J.M. 1986. *The use of an enzymatic technique to predict digestibility, metabolisable and net energy of compound feedstuffs for ruminants*. Animal Feed Science and Technology 14:203-214.
- [16] Vintilă, M., 1989. *O metodă eficientă de conservare a nutrețurilor*. Revista de creșterea animalelor, 7.
- [17] Demarquilly, C., Andrieu, J., 1988. *Les fourrages*. In R. JARRIGE, *Alimentation des bovins, ovins & caprins*. Paris: INRA Editions, 315-355.
- [18] Williams, P., Sobering, D., 1992. In *Near Infrared Spectroscopy: Bridging the gap between data analysis and NIR applications*. Proceedings of the 5th ICNIRS (pp. 441-446). Haugesund, Norway: Ellis Horwood Ltd, Chichester.
- [19] Sheaffer, C. C., Martin, N. P., Lamb, J. F., Cuomo, G. R., Jewett, J. G., 2000. *Leaf and Stem Properties of Alfalfa Entries*. Agronomy Journal, 733-739.
- [20] Velasco, L., Fernandez-Martinez, J., De Haro, A., 1998. *Application of near-infrared reflectance spectroscopy to estimate the bulk density of Ethiopian mustard seeds*. Journal of the Science of Food and Agriculture, 77 (3), 312-318.
- [21] Iancheva, N., Steingass, H., Todorov, N., Pavlov, D., 1999. *A comparison of in vitro rumen fluid and enzymatic methods to predict digestibility and energy value of grass and alfalfa hay*. Animal Feed Science and Technology, 81, 333-344.
- [22] Yu, P., Christensen, D. A., McKinnon, J. J., Markert, J. D. 2003. *Effect of variety and maturity stage on chemical composition, carbohydrate and protein subfractions, in vitro rumen degradability and crude energys of timothy and alfalfa*. Canadian Journal of Animal Science, 279-290.
- [23] INRA, 2007. *Alimentation des bovins, ovins et caprins. Besoins des animaux. Valeur des aliments*. Tables INRA 2007. Editions Quæ, Paris, France, 307p.
- [24] Homolka, P., Koukolova, V., Němec, Z., Mudřik, Z., Hučko, B., Sales, J., 2008. *Amino acid contents and intestinal digestibility of lucerne in ruminants as influenced by growth stage*. Czech Journal of Animal Science, 499-505.
- [25] Stanacev, V., Dukic, D., Kovcin, S., Drinic, M., Puva, N., Stanacev, V., 2010. *Nutritive value of the genetically divergent genotypes of lucerne (Medicago*

sativa L.). African Journal of Agricultural Research, 5 (11), 1284-1287.

[27]Heinriches, D., 1970. *Variation of chemical constituents within and between alfalfa populations*. In M. Norman, 11th Proc. Int. Grassl. Congr. (pp. 267-270). St. Lucia, Queensland: University of Queensland Press.

[28]Babinec, J., Kozova, Z., Strakova, E., Suchy, P., 2001. *The variance of the amino acids in some lucerne (Medicago sativa L.) populations*. In I. Delgado, & J.

Lloveras, *Quality in lucerne and medics for animal production* (pp. 235-239). Zaragoza (Spain): CIHEAM.

[28]Brognna, N., Pacchioli, M. T., Immovilli, A., Ruozzi, F., Ward, R., Formigoni, A., 2009. *The use of near-infrared reflectance spectroscopy (NIRS) in the prediction of chemical composition and in vitro neutral detergent fiber (NDF) digestibility of Italian alfalfa hay*. Italian Journal of Animal Science, 271-273.



ISSN 2285 – 5750
ISSN-L 2285 – 5750